

On solutions to the non-Abelian Hirota-Miwa equation and its continuum limits

C.X. Li¹, J.J.C. Nimmo² and K.M. Tamizhmani³

¹School of Mathematical Sciences,
Capital Normal University
Beijing 100048, CHINA

²Department of Mathematics,
University of Glasgow
Glasgow G12 8QW, UK

³Department of Mathematics,
Pondicherry University
Kalapet, 605014 Pondicherry, India

Abstract

In this paper, we construct grammian-like quasideterminant solutions of a non-Abelian Hirota-Miwa equation. Through continuum limits of this non-Abelian Hirota-Miwa equation and its quasideterminant solutions, we construct a cascade of noncommutative differential-difference equations ending with the noncommutative KP equation. For each of these systems the quasideterminant solutions are constructed as well.

1 Introduction

Recently, noncommutative versions of some well-known soliton equations have been extensively studied. These systems include the KP equation, the KdV equation, the Hirota-Miwa equation, the modified KP equation and the two-dimensional Toda lattice equation [1–14]. Generally speaking, determinants are ubiquitous as solutions of commutative integrable systems while quasideterminants are often solutions of noncommutative integrable systems [15–17]. In many cases, the noncommutative version of the equation is obtained as the compatibility of the same Lax pair for the commutative version. However, in the noncommutative case, the assumption that the coefficients in the Lax pair commute is relaxed.

This paper is concerned with the non-Abelian version of the Hirota-Miwa equation introduced in [10]. Given the system of linear equations

$$a_i^{-1}T_i(\phi) - a_j^{-1}T_j(\phi) = U_{ij}\phi, \quad i, j \in \{1, 2, 3\}, \quad (1.1)$$

where ϕ and U_{ij} belong to an associative algebra $\mathcal{A}$ and depend on variables n_1, n_2, n_3 , T_i denotes the shift operator in variable n_i , defined by $T_i(X) = X(n_i + 1)$ and a_i are nonzero scalar constants, called lattice parameters. We will also use the shorthand subscript notation $X_{,i} = T_i(X)$. This linear system is compatible if and only if

$$U_{ij} + U_{jk} + U_{ki} = 0, \quad (1.2)$$

$$T_k(U_{ij}) + T_i(U_{jk}) + T_j(U_{ki}) = 0, \quad (1.3)$$

$$T_k(U_{ij})U_{ki} = T_j(U_{ki})U_{ij}, \quad (1.4)$$

for each $i, j, k \in \{1, 2, 3\}$. In particular, these give $U_{ii} = 0$ and $U_{ij} = -U_{ji}$ for $i, j \in \{1, 2, 3\}$. Equations (1.2)–(1.4) are referred to as the non-Abelian Hirota-Miwa equation [10, 12] since in the commutative case if we make the ansatz

$$U_{ij} = (a_i^{-1} - a_j^{-1}) \frac{T_{ij}(\tau)\tau}{T_i(\tau)T_j(\tau)}, \quad (1.5)$$

where $T_{ij}(\tau)$ denotes $T_i(T_j(\tau))$, then all of the equations in (1.2)–(1.4) reduce to just one equation, the well-known Hirota-Miwa equation [18, 19]

$$a_1(a_2 - a_3)T_1(\tau)T_{23}(\tau) + a_2(a_3 - a_1)T_2(\tau)T_{31}(\tau) + a_3(a_1 - a_2)T_3(\tau)T_{12}(\tau) = 0. \quad (1.6)$$

In the commutative case, it is well known that the Miwa transformation [19]

$$x_k = \sum_{i=1}^{\infty} \frac{a_i^k}{k} n_i,$$

provides the link between the discrete variables n_i and the continuous variables x_k in the KP hierarchy. Using this transformation, we can view $\tau = \tau(n_1, n_2, n_3, \dots; x_1, x_2, x_3, \dots)$ as a function of both discrete and continuous variables such that

$$T_i(\tau) = \tau|_{n_i \rightarrow n_i+1} = \tau|_{x_1 \rightarrow x_1+a_i, x_2 \rightarrow x_2+a_i^2/2, x_3 \rightarrow x_3+a_i^3/3, \dots}, \quad (1.7)$$

and the continuum limits, as lattice parameters $a_i \rightarrow 0$, are obtained by using Taylor's theorem. In this way we have, for example,

$$T_i(\tau) = \tau + a_i \tau_x + \frac{1}{2} a_i^2 (\tau_{xx} + \tau_y) + \frac{1}{6} a_i^3 (\tau_{xxx} + 3\tau_{xy} + 2\tau_t) + O(a_i^4),$$

where, for a shorter notation, we have written x_1, x_2, x_3 as x, y, t respectively.

Taking successive continuum limits one obtains a cascade of differential-difference equations starting from the Hirota-Miwa equation and ending with the KP equation. To be precise, as $a_3 \rightarrow 0$ the leading order term in (1.6) is at $O(a_3)$ and is

$$(a_1 - a_2)(T_1(\tau)T_2(\tau) - T_{12}(\tau)\tau) + a_1 a_2 D_x T_1(\tau) \cdot T_2(\tau) = 0, \quad (1.8)$$

which we call the $\Delta^2 \partial \text{KP}$ equation. Next, as $a_2 \rightarrow 0$ the leading order term in (1.8) is at $O(a_2^2)$ and is

$$(a_1 D_y + a_1 D_x^2 - 2D_x) T_1(\tau) \cdot \tau = 0, \quad (1.9)$$

which is nothing but the differential-difference KP equation. Here we denote (1.9) as the $\Delta \partial^2 \text{KP}$ equation to stress that it has one discrete variable and two continuous ones. Finally, as $a_1 \rightarrow 0$ the leading order term in (1.8) is at $O(a_1^3)$ and is

$$(D_x^4 + 3D_y^2 - 4D_x D_t) \tau \cdot \tau = 0, \quad (1.10)$$

the Hirota form of the KP equation. The whole procedure is illustrated in Figure 1.




Figure 1: Continuum limits

The main aim of this paper is to show that the same cascade of continuum limits also exists in the non-Abelian case and thereby we construct noncommutative versions of the differential-difference KP equations (1.8) and (1.9). We also give solutions of these systems in terms of quasideterminants.

The paper is organized as follows. In Section 2, we review results from [10] on the Daroux transformations for the non-Abelian Hirota-Miwa equation and consider the binary Darboux transformation and the quasigrammian solutions of (1.2)–(1.4) that it constructs. Using the ansatz that is suggested by this binary Darboux transformation, in Section 3, we first reformulate the results presented in a way which facilitates

taking continuum limits. Then, by taking the successive continuum limits described above we first obtain a noncommutative version of (1.8). Quasiwronskian solutions and quasigrammian solutions for this system are also obtained. Then a second continuum limit yields a noncommutative version of (1.9) and its quasiwronskian and quasigrammian solutions are presented. Finally we show that the continuum limit of this noncommutative differential-difference KP is nothing but the noncommutative KP equation. Conclusions and discussions are given in Section 4.

2 The non-Abelian Hirota-Miwa equation

In this section, we will present quasigrammian solutions to the non-abelian Hirota-Miwa equation (1.2)–(1.4) constructed by using binary Darboux transformations.

Let us first review some results on the non-Abelian Hirota-Miwa equation. Quasicasoratian solutions were constructed for the non-Abelian Hirota-Miwa equation by repeating Darboux transformations in [10,12]. The linear system (1.1) considered here is a generalisation of the one used in [10,12] but it is not difficult to prove that (1.2)–(1.4) has the Darboux transformation

$$\tilde{\phi} = a_i^{-1}(T_i(\phi) - T_i(\theta)\theta^{-1}\phi), \quad (2.1)$$

$$\tilde{U}_{ij} = T_j(T_i(\theta)\theta^{-1})U_{ij}\theta T_i(\theta)^{-1}, \quad i, j \in \{1, 2, 3\} \quad (2.2)$$

where θ is a particular solutions of the linear system (1.1). This suggests the ansatz $U_{ij} = T_j(C_i)U_{i,j}^0 C_i^{-1}$, where $U_{i,j}^0$ is a seed solution. Then iterating the Darboux transformation gives casoratian-type quasideterminant solutions

$$C_i[n] = \begin{vmatrix} \theta & 1 \\ T_i(\theta) & 0 \\ \vdots & \vdots \\ T_i^{n-1}(\theta) & 0 \\ T_i^n(\theta) & \boxed{0} \end{vmatrix} C_i, \quad (2.3)$$

where $\theta = (\theta_1, \dots, \theta_N)$ is a vector solution of (1.1). In the next section we will rewrite this Darboux transformation using a different parametrization and obtain a different casoratian-type solution.

To construct binary Darboux transformations for (1.2)–(1.4), we must define an appropriate adjoint system. The formal adjoint of the linear system (1.1) has the form

$$a_i^{-1}T_i^{-1}(\psi) - a_j^{-1}T_j^{-1}(\psi) = V_{ij}\psi, \quad i, j \in \{1, 2, 3\}.$$

and we will choose V_{ij} in order that the compatibility condition of this adjoint system is equivalent to (1.2)–(1.4). The compatibility condition for this system of linear equations can be deduced from (1.2)–(1.4) by simply changing T_i to T_i^{-1} and U_{ij} to V_{ij} in this system and we get

$$V_{ij} + V_{jk} + V_{ki} = 0, \quad (2.4)$$

$$T_k^{-1}(V_{ij}) + T_i^{-1}(V_{jk}) + T_j^{-1}(V_{ki}) = 0, \quad (2.5)$$

$$T_k^{-1}(V_{ij})V_{ki} = T_j^{-1}(V_{ki})V_{ij}. \quad (2.6)$$

This is equivalent to (1.2)–(1.4) when $T_{ij}(V_{ij}) = U_{ij}^*$, where $*$ is an involutive antiautomorphism defined on $\mathcal{A}$ satisfying $(AB)^* = B^*A^*$ and $(A^*)^* = A$. Thus we obtain the adjoint linear system in its final form

$$a_i^{-1}T_j(\psi) - a_j^{-1}T_i(\psi) = U_{ij}^*T_{ij}(\psi). \quad (2.7)$$

An eigenfunction potential $\Omega(\phi, \psi)$ may be defined by

$$\Delta_i(\Omega) = T_i(\psi^*)\phi, \quad (2.8)$$

for all i , where $\Delta_i = a_i^{-1}(T_i - 1)$. It is straightforward to show that the exactness condition $(\Omega_{,i})_{,j} = (\Omega_{,j})_{,i}$ when ϕ and ψ satisfy (1.1) and (2.7) respectively. Following the standard construction, a binary Darboux

transformation is then defined by

$$\tilde{\phi} = \phi - \theta \Omega(\theta, \rho)^{-1} \Omega(\phi, \rho), \quad (2.9)$$

$$\tilde{\psi} = \psi - \rho(\Omega(\theta, \rho)^*)^{-1} \Omega(\theta, \psi)^*, \quad (2.10)$$

$$\tilde{U}_{ij} = U_{ij} - (T_i - T_j)(\theta \Omega(\theta, \rho)^{-1} \rho^*), \quad (2.11)$$

where θ and ρ are particular solutions of the linear system (1.1) and its adjoint (2.7).

There is also a ‘vector’ form of this transformation, equivalent to an iterated transformation which gives solutions

$$U_{ij} = U_{ij}^0 - (T_i - T_j)(\theta \Omega(\theta, P)^{-1} P^*), \quad (2.12)$$

where U_{ij}^0 is a seed solution of (1.2)–(1.4), θ is as defined above and $P = (\rho_1, \dots, \rho_N)$ is a vector solution of the adjoint linear problem (2.7) with potential $U_{i,j} = U_{ij}^0$. This solution may also be expressed as a quasideterminant

$$U_{ij} = U_{ij}^0 + (T_i - T_j) \left| \begin{array}{c} \Omega(\theta, P) \\ \theta \end{array} \right| \frac{P^*}{\boxed{0}}. \quad (2.13)$$

This result can be proved by induction but here we choose to verify the solution by direct substitution into (1.2)–(1.4). We write

$$U_{ij} = U_{ij}^0 + G_{,i} - G_{,j}, \quad (2.14)$$

where

$$G = \left| \begin{array}{c} \Omega(\theta, P) \\ \theta \end{array} \right| \frac{P^*}{\boxed{0}} = -\theta \Omega^{-1} P^*. \quad (2.15)$$

Observe that (1.2), (1.3) are identically satisfied by the ansatz (2.14), irrespective of the form of G , and all nontrivial versions of (1.4), in which i, j and k are distinct, give the same equation

$$G_{,12}(U_{12}^0 + G_{,1} - G_{,2}) + G_{,23}(U_{23}^0 + G_{,2} - G_{,3}) + G_{,31}(U_{31}^0 + G_{,3} - G_{,1}) + (U_{23}^0 G)_{,1} + (U_{31}^0 G)_{,2} + (U_{12}^0 G)_{,3} = 0. \quad (2.16)$$

Using (2.8), we obtain for any i, j, k

$$G_{,ik} G_{,k} = a_i^{-1} \theta_{,ik} (\Omega_{,k}^{-1} - \Omega_{,ik}^{-1}) P_{,k}^*.$$

Then substituting these expressions into (2.16), and making use of the linear equations (1.1) and (2.7), it is straightforward to show that (2.16) is identically satisfied. This completes the verification of the solutions (2.13).

3 Continuum limits of the non-Abelian Hirota-Miwa equation

In this section, we will find the proper continuum limits of the non-Abelian Hirota-Miwa equation as the lattice parameters tend to zero. As well as constructing the nonlinear equations, the corresponding linear equations (Lax pairs), Darboux and binary Darboux transformations and solutions will be obtained also in the same limit.

The first step is to reformulate the results for the non-Abelian Hirota-Miwa equation in a way that facilitates these continuum limits. In particular, we reexpress many terms using the forward and backward difference operators $\Delta_i := a_i^{-1}(T_i - 1)$ and $\nabla_i := a_i^{-1}(1 - T_i^{-1})$. We make an ansatz of the form (2.14) suggested by the binary Darboux transformation, in which U_{ij}^0 is any seed solution of (1.2)–(1.4). In fact, in order to successfully balance leading order terms, it turns out to be necessary to choose the seed solution in the form $U_{ij}^0 = a_i^{-1} - a_j^{-1}$. Thus we write the solution of (1.2)–(1.4) as

$$U_{ij} = a_i^{-1} - a_j^{-1} + T_i G - T_j G. \quad (3.1)$$

Under this assumption, the system (1.2)–(1.4) reduces to the single equation

$$\begin{aligned}
& a_1 a_2 (\Delta_1 \Delta_2 G (a_1 \Delta_1 G - a_2 \Delta_2 G) - [\Delta_1 G, \Delta_2 G]) \\
& + a_2 a_3 (\Delta_2 \Delta_3 G (a_2 \Delta_2 G - a_3 \Delta_3 G) - [\Delta_2 G, \Delta_3 G]) \\
& + a_3 a_1 (\Delta_3 \Delta_1 G (a_3 \Delta_3 G - a_1 \Delta_1 G) - [\Delta_3 G, \Delta_1 G]) = \\
& (a_1 - a_2) \Delta_1 \Delta_2 G + (a_2 - a_3) \Delta_2 \Delta_3 G + (a_3 - a_1) \Delta_3 \Delta_1 G.
\end{aligned} \tag{3.2}$$

and its linear system and adjoint linear system may be written as

$$\Delta_i \phi - \Delta_j \phi = (a_i \Delta_i G - a_j \Delta_j G) \phi, \tag{3.3}$$

and

$$-\nabla_i \psi + \nabla_j \psi = (a_i \nabla_i G^* - a_j \nabla_j G^*) \psi, \tag{3.4}$$

respectively.

For the purpose of taking continuum limits, it is more convenient to reformulate the Darboux transformation (2.1)–(2.2) using the parametrization (2.14) given by the binary Darboux transformation. Doing this we find that

$$\tilde{\phi} = \Delta_i \phi - (\Delta_i \theta) \theta^{-1} \phi, \tag{3.5}$$

$$\tilde{G} = T_i G - (\Delta_i \theta) \theta^{-1}. \tag{3.6}$$

Superficially, these expressions seem strange as the LHS depends on index i but the RHS does not. It is however easy to check using the linear problem (3.3) that the RHS in each case is independent of the choice of i . Iterating the Darboux transformation in this form again gives quasicasoratian solutions but different from the ones given in (2.3),

$$G[n] = T_i^n G + \begin{vmatrix} \theta & 0 \\ \vdots & \vdots \\ \Delta_i^{n-2} \theta & 0 \\ \Delta_i^{n-1} \theta & 1 \\ \Delta_i^n \theta & \boxed{0} \end{vmatrix}, \tag{3.7}$$

where θ is as in (2.3). Again, it may be shown that the RHS of (3.7) is independent of the choice of i .

3.1 Noncommutative $\Delta^2 \partial$ KP equation

To take the first continuum limit, let $a_3 \rightarrow 0$ and note that, for example, $\Delta_3 G = G_x + O(a_3)$.

Nonlinear equation The leading order terms in (3.2) give

$$a_1 a_2 (\Delta_1 \Delta_2 G (a_1 \Delta_1 G - a_2 \Delta_2 G) - [\Delta_1 G, \Delta_2 G]) = (a_1 - a_2) \Delta_1 \Delta_2 G + a_2 \Delta_2 G_x - a_1 \Delta_1 G_x, \tag{3.8}$$

which we call the noncommutative $\Delta^2 \partial$ KP equation.

Lax pair The leading order terms in (3.3) for give the Lax pair

$$\Delta_1 \phi = \phi_x + a_1 (\Delta_1 G) \phi \tag{3.9}$$

$$\Delta_2 \phi = \phi_x + a_2 (\Delta_2 G) \phi. \tag{3.10}$$

Similarly, the leading order terms in (3.4) give the adjoint form

$$\nabla_1 \psi = \psi_x - a_1 (\nabla_1 G^*) \psi, \tag{3.11}$$

$$\nabla_2 \psi = \psi_x - a_2 (\nabla_2 G^*) \psi. \tag{3.12}$$

The compatibility conditions of both (3.9)–(3.10) and (3.11)–(3.12) are (3.8).

Darboux transformation Similarly from (3.5)–(3.6), we get the Darboux transformation for (3.9)–(3.10), either expressed in terms of the x -derivative or difference operators Δ_i

$$\tilde{\phi} = \phi_x - \theta_x \theta^{-1} \phi = \Delta_i \phi - (\Delta_i \theta) \theta^{-1} \phi, \quad (3.13)$$

$$\tilde{G} = G - \theta_x \theta^{-1} = T_i G - (\Delta_i \theta) \theta^{-1}, \quad (3.14)$$

for $i = 1, 2$.

Binary Darboux transformation It is clear that the binary Darboux transformation always keeps the same form as (2.9)–(2.11) whatever the continuum limits are. The only difference is the definition of the eigenfunction potential Ω . Here and hereafter, we will only give the definition of Ω . The leading order terms in (2.8) give the eigenfunction potential

$$\Omega_x = \psi^* \phi, \quad (3.15)$$

$$\Delta_i \Omega = T_i(\psi^*) \phi, \quad i = 1, 2, \quad (3.16)$$

Solutions From the continuum limit of the quasicasoratian solutions (3.7) for $i = 3$, we get an alternative quasiwronskian expression and so the Darboux transformations give solutions

$$G[n] = G + \begin{vmatrix} \theta & 0 \\ \vdots & \vdots \\ \theta^{(n-2)} & 0 \\ \theta^{(n-1)} & 1 \\ \theta^{(n)} & \boxed{0} \end{vmatrix} = T_i^n G + \begin{vmatrix} \theta & 0 \\ \vdots & \vdots \\ \Delta_i^{n-2} \theta & 0 \\ \Delta_i^{n-1} \theta & 1 \\ \Delta_i^n \theta & \boxed{0} \end{vmatrix}, \quad (3.17)$$

for $i = 1, 2$. The quasigrammian solutions are given by (2.15) with Ω defined by (3.15)–(3.16).

3.2 Noncommutative $\Delta \partial^2$ KP equation

To take the second continuum limit, let $a_2 \rightarrow 0$ and note that, for example, $\Delta_2 G = G_x + \frac{1}{2} a_2 (G_{xx} + G_y) + O(a_2^2)$ and $\nabla_2 G = G_x + \frac{1}{2} a_2 (G_y - G_{xx}) + O(a_2^2)$.

Nonlinear equation The leading order terms in (3.8) give the noncommutative differential-difference version of the potential KP equation

$$a_1^2 \Delta_1 G_x \Delta_1 G - a_1 [\Delta_1 G, G_x] = -\Delta_1 G_x + \frac{1}{2} a_1 (\Delta_1 G_{xx} + \Delta_1 G_y) + G_{xx}. \quad (3.18)$$

Lax pair The leading order terms in (3.9) and (3.10) give the Lax pair

$$\phi_y + \phi_{xx} = 2G_x \phi, \quad (3.19)$$

$$\Delta_1 \phi = \phi_x + a_1 (\Delta_1 G) \phi. \quad (3.20)$$

The leading order terms in (3.11)–(3.12) give the adjoint Lax pair

$$\psi_y - \psi_{xx} + 2G_x^* \psi = 0, \quad (3.21)$$

$$\nabla_1 \psi = \psi_x - a_1 (\nabla_1 G^*) \psi. \quad (3.22)$$

The compatibility conditions of both (3.19)–(3.20) and (3.21)–(3.22) are (3.18).

Darboux transformation Similarly, from (3.13)–(3.14), we get the Darboux transformation for (3.19)–(3.20), either expressed in terms of x -derivative or the difference operator Δ_1

$$\tilde{\phi} = \phi_x - \theta_x \theta^{-1} \phi = \Delta_1 \phi - (\Delta_1 \theta) \theta^{-1} \phi, \quad (3.23)$$

$$\tilde{G} = G - \theta_x \theta^{-1}. \quad (3.24)$$

Binary Darboux transformation Considering the expansion of (3.16) for $i = 2$ as $a_2 \rightarrow 0$ gives

$$\Omega_x + \frac{1}{2}a_2(\Omega_{xx} + \Omega_y) = (\psi^* + a_2\psi_x^*)\phi + O(a_2^2).$$

From this it follows that $\Omega_y = 2\psi_x^*\phi - \Omega_{xx}$ and so the eigenfunction potential Ω satisfies

$$\Omega_x = \psi^*\phi, \quad (3.25)$$

$$\Omega_y = \psi_x^*\theta - \psi^*\theta_x, \quad (3.26)$$

$$\Delta_1\Omega = (T_1\psi)^*\phi. \quad (3.27)$$

Solutions The Darboux transformations give solutions

$$G[n] = G + \begin{vmatrix} \theta & 0 \\ \vdots & \vdots \\ \theta^{(n-2)} & 0 \\ \theta^{(n-1)} & 1 \\ \theta^{(n)} & \boxed{0} \end{vmatrix} = T_1^n G + \begin{vmatrix} \theta & 0 \\ \vdots & \vdots \\ \Delta_1^{n-2}\theta & 0 \\ \Delta_1^{n-1}\theta & 1 \\ \Delta_1^n\theta & \boxed{0} \end{vmatrix}, \quad (3.28)$$

and the quasigrammian solutions are given by (2.15) with Ω defined by (3.25)–(3.27).

3.3 Noncommutative KP equation

To take the final continuum limit, let $a_1 \rightarrow 0$ and note that here $\Delta_1 G = G_x + \frac{1}{2}a_1(G_y + G_{xx}) + \frac{1}{6}a_1^2(G_{xxx} + 3G_{xy} + 2G_t) + O(a_1^3)$ and $\nabla_1 G = G_x + \frac{1}{2}a_1(G_y - G_{xx}) + \frac{1}{6}a_1^2(2G_t - 3G_{xy} + G_{xxx}) + O(a_1^3)$.

Nonlinear equation The leading order terms in (3.18) give the noncommutative potential KP equation

$$(-4G_t - 6G_x^2 + G_{xxx})_x + 3G_{yy} = 6[G_x, G_y]. \quad (3.29)$$

Lax pair The leading order terms in (3.19)–(3.20) give the Lax pair

$$\phi_y = -\phi_{xx} + 2G_x\phi, \quad (3.30)$$

$$\phi_t = \phi_{xxx} - 3G_x\phi_x + \frac{3}{2}(G_y - G_{xx})\phi. \quad (3.31)$$

Similarly, the leading order terms in (3.21)–(3.22) give the adjoint form

$$\psi_y = \psi_{xx} - 2G_x^*\psi, \quad (3.32)$$

$$\psi_t = \psi_{xxx} - 3G_x^*\psi_x - \frac{3}{2}(G_y^* + G_{xx}^*)\psi. \quad (3.33)$$

The compatibility conditions of both (3.30)–(3.31) and (3.32)–(3.33) are (3.29).

Darboux transformation The leading orders in (3.23)–(3.24) give the Darboux transformation for (3.29)

$$\tilde{\phi} = \phi_x - \theta_x\theta^{-1}\phi, \quad (3.34)$$

$$\tilde{G} = G - \theta_x\theta^{-1}. \quad (3.35)$$

Binary Darboux transformation Considering the expansion of (3.16) for $i = 2$ as $a_1 \rightarrow 0$ gives

$$\Omega_x + \frac{1}{2}a_1(\Omega_{xx} + \Omega_y) + \frac{1}{6}a_1^2(\Omega_{xxx} + 3\Omega_{xy} + 2\Omega_t) = (\psi^* + a_1\psi_x^* + \frac{1}{2}a_1^2(\psi_{xx}^* + \psi_y^*))\phi + O(a_1^3).$$

From this it follows that $2\Omega_t = 3(\psi_{xx}^* + \psi_y^*)\phi - \Omega_{xxx} - 3\Omega_{xy}$ and so the eigenfunction potential Ω satisfies

$$\Omega_x = \psi^*\phi, \quad (3.36)$$

$$\Omega_y = \psi_x^*\phi - \psi^*\phi_x, \quad (3.37)$$

$$\Omega_t = \psi_{xx}^*\phi - \psi_x^*\phi_x + \psi^*\phi_{xx} - 3\psi^*G_x\phi. \quad (3.38)$$

Remark. In [11], the following noncommutative KP equation is considered

$$(U_t + 3U_x^2 + U_{xxx})_x + 3U_{yy} = 3[U_x, U_y]. \quad (3.39)$$

Its quasideterminant solutions are obtained through Darboux transformation and binary Darboux transformation. In fact, using the scaling transformations $y \rightarrow -y$, $t \rightarrow -4t$ and writing in terms of $U = -2G$, (3.29) can be transformed into (3.39).

Solutions Finally, we recover the quasideterminant solutions of the noncommutative KP equation, either quasiwronskian [11, 16] or quasigrammian [11]

$$U[n] = U - 2 \begin{vmatrix} \theta & 0 \\ \vdots & \vdots \\ \theta^{(n-2)} & 0 \\ \theta^{(n-1)} & 1 \\ \theta^{(n)} & \boxed{0} \end{vmatrix}, \text{ or } U[n] = U - 2 \begin{vmatrix} \Omega(\theta, P) & P^* \\ \theta & \boxed{0} \end{vmatrix}.$$

4 Conclusions and discussion

In this paper, we first obtained the quasigrammian solutions of the non-Abelian Hirota-Miwa equation by using an iterated binary Darboux transformation. The binary Darboux transformation leads to an ansatz (2.14) for U_{ij} which reduces the non-Abelian Hirota-Miwa equations to a single equation symmetric in the three discrete variables n_i . Then by considering the continuum limits of this equation and its Darboux transformations and binary Darboux transformations, we obtain a cascade of noncommutative differential-difference equations and their quasideterminant solutions.

There is a second parametrization, $U_{ij} = (a_i^{-1} - a_j^{-1})T_j(C_i)C_i^{-1}$ which can be seen from the quasiwronskian solution constructed by its Darboux transformation. A natural question is ‘Can we obtain continuum limits of the non-Abelian Hirota-Miwa equation under the second parametrization?’

Using this parametrization the linear equations are

$$\Delta_i \phi - \Delta_j \phi = a_j(a_i^{-1} - a_j^{-1})\Delta_j(C_i)C_i^{-1}\phi, \quad i, j \in \{1, 2, 3\},$$

and in the limit $a_3 \rightarrow 0$, this system becomes

$$\begin{aligned} \Delta_1 \phi - \Delta_2 \phi &= a_2(a_1^{-1} - a_2^{-1})\Delta_2(C_1)C_1^{-1}\phi, \\ \Delta_1 \phi &= \phi_x - C_{1x}C_1^{-1}\phi, \\ \Delta_2 \phi &= \phi_x - C_{2x}C_2^{-1}\phi. \end{aligned}$$

The algebraic compatibility of these equations gives

$$a_2(a_1 - a_2)\Delta_2(C_1)C_1^{-1} + a_1a_2(C_{2x}C_2^{-1} - C_{1x}C_1^{-1}) = 0, \quad (4.1)$$

and it may be shown also that the only other compatibility condition, $\phi_{,12} = \phi_{,21}$ is identically satisfied provided (4.1) is satisfied. This, as expected, agrees with the leading order terms in the continuum limit of $U_{12} + U_{23} - U_{13} = 0$.

Now let $u := C_{1x}C_1^{-1}$ and $v := C_{2x}C_2^{-1}$, then (4.1) can be rewritten as

$$(u - v)_x + (a_1^{-1} - a_2^{-1})a_2\Delta_2(u) + a_2\Delta_2(u)v - a_1\Delta_1(v)u + [u, v] = 0. \quad (4.2)$$

On the other hand, exactly the same equation arises if we write $u = -a_1\Delta_1(G)$ and $v = -a_2\Delta_2(G)$ in the noncommutative $\Delta^2\partial$ KP equation (3.8). Thus we see that in the first continuum limit the second parametrization gives a system equivalent to that obtained from the first. However, it is not at all clear how to take the second continuum limit, $a_2 \rightarrow 0$. For this reason we have not pursued the second parametrization any further.

Acknowledgement

This work was supported in part by a Royal Society China Fellowship and the National Natural Science Foundation of China (grant no. 10601028). K.M. Tamizhmani wishes to acknowledge the NBHM for the project and J.J.C. Nimmo wishes to acknowledge the support of a Royal Society of London International Outgoing Short Visit award.

References

- [1] B.A. Kupershmidt, KP or mKP: Noncommutative Mathematics of Lagrangian, Hamiltonian, and Integrable systems. Mathematical Surveys and Monographs (American Mathematical Society, New York), 78, 2000.
- [2] L.D. Paniak, Exact noncommutative KP and KdV multi-solitons, arXiv:hep-th/0105185 (2001).
- [3] M. Sakakibara, J. Phys. A 37 (2004) L599-L604.
- [4] N. Wang, M. Wadati, J. Phys. Soc. Jpn 72 (2003) 1366-1373.
- [5] N. Wang, M. Wadati, J. Phys. Soc. Jpn 72 (2003) 1881-1888.
- [6] N. Wang, M. Wadati, J. Phys. Soc. Jpn 73 (2004) 1689-1698.
- [7] M. Hamanaka, Noncommutative solitons and D-branes. PhD thesis. arXiv:hep-th/0303256 (2003).
- [8] M. Hamanaka, Nucl. Phys. B 316 (2003) 77-83.
- [9] A. Dimakis, Müller-Hoissen, J. Phys. A 38 (2005) 5453-5505.
- [10] J.J.C. Nimmo, J. Phys. A 39 (2006) 5053-5065.
- [11] C.R. Gilson, J.J.C. Nimmo, J. Phys. A 40 (2007) 3839-3850.
- [12] C.R. Gilson, J.J.C. Nimmo, Y. Ohta, J. Phys. A: Math. Theor. 40 (2007) 12607-12617.
- [13] C.R. Gilson, J.J.C. Nimmo, C.M. Sooman, J. Phys. A: Math. Theor. 41 (2008) 085202.
- [14] C.X. Li, J.J.C. Nimmo, Proc. R. Soc. A 464 (2008) 951-966.
- [15] I.M. Gelfand, V.S. Retakh, Funkt. Anal. Prilozhen. 25 (1991) 13-25.
- [16] P. Etingof, I.M. Gelfand, V.S. Retakh, Math. Res. Lett. 5(1998) 1-12.
- [17] I.M. Gelfand, S. Gelfand, V.M. Retakh, R.L. Wilson, Adv. Math. 193 (2005) 56-141.
- [18] R. Hirota, J. Phys. Soc. Japan 50 (1982) 3785-91.
- [19] T. Miwa, Proc. Japan Acad. A 58 (1982) 9-12.